



Republic of the Philippines
VISAYAS STATE UNIVERSITY
 Visca, Baybay City, Leyte

PURCHASE REQUEST

PR No.: **GF-2022-08-01154**Date: **08-23-2022**Dept./Office: **DVCS**

Category: **Medical &
 Dental Supplies &
 Equipment**

Section/End-User: **Shebelle A. Cueva**Funding Source: **General Fund - MOOE**Project Title/Code: **CVM/ Extension (VetMends)**

Item #	Item Description	Unit	Qty	Unit Cost	PAR/ICS	Total Cost
1	Amoxicillin trihydrate 150 mg/mL	btl	1	900.00		900.00

Specification:

Amoxicillin trihydrate 150 mg/mL (100 mL/bottle)

suspension for Injection (IM,SC)

Antibacterial

FOR VETERINARY USE ONLY

FORMULATION:

each mL contains:

Amoxicillin trihydrate equivalent to amoxicillin 150 mg

INDICATIONS:

long acting suspension used for the treatment of localised infections, respiratory infections, and secondary bacterial infections caused by organisms sensitive to amoxicillin.

2	Atropine sulfate 1mg/ml/amp	Ampule	1	33.00		33.00
3	Disposable syringewith needle, 3cc G-23	Box of 100's	2	650.00		1,300.00
4	Epinephrine hydrochloride 1mg/ml/ampule	Box of 10's	1	800.00		800.00
5	Gauze Pad 4x4x8 Ply (sterile)	box of 20 packs	3	400.00		1,200.00
6	gloves surgical, sterile, powder free 6.5	box of 50 pairs	1	900.00		900.00

7	gloves surgical, sterile, powder free 7.5	box of 50pairs	1	900.00		900.00
8	Hydrogen peroxide 3% solution 1L	btl	2	110.00		220.00
9	Povidone Iodine wound solution 7.5%/ gal	gallon	1	725.00		725.00
10	tolfenamic acid/anti-inflammatory drug	btl	1	2,900.00		2,900.00

Specification:

Tolfenamic acid 40 mg/mL (100 mL/bottle)

Solution for injection

FORMULATION:

Sterile aqueous solution containing 4.0% w/v tolfenamic acid as active ingredient and 1.04% w/v benzyl alcohol as preservative.

INDICATION:

is a non steroidal anti-inflammatory of the fenamate class.

Its active principle, Tolfenamic acid, plays a double role in prostaglandins:

- inhibition of prostaglandin synthesis from arachidonic acid.
- ?• inhibition of tissue response to prostaglandins by occupation of receptor sites.

Tolfenamic acid is a potent pain reliever.

DIRECTION:

Restraints: DO NOT inject cattle other than into muscle tissue high on the side of the neck.

Injection of product into muscles other than as described is likely to result in residues in meat above the MRL or ESI.

IV administration is also available for Tolfedine® CS.

Cattle






Pneumonia: 1 ml/20 kg by intramuscular injection. Treatment may be repeated once only after 48 hours.

Mastitis: 1 ml per 10 kg as a single intravenous injection.

Pigs

Metritis Mastitis Agalactia: 1 ml/20 kg bw as a single intramuscular injection into the rump.

11	Tranexamic acid 100mg/ml/5ml ampule	Box of 10's	1	2,000.00		2,000.00
12	Tuberculin syringe	box of 50's	2	900.00		1,800.00
13	Zoletil	btl	3	1,320.00		3,960.00

TOTAL				17,638.00	
Purpose: Veterinary Medical Mission Supplies					
Checked by:  PHOEBE LYNN B. CALUNGSOD			Funds Available:  ALICIA M. FLORES <i>EXT 77</i>		
TWG - Medical & Dental Supplies & Equipment			HEAD, BUDGET OFFICE		
Signature: Printed Name: Designation:	Requested by:  SHEBELLE A. CUEVA		Noted by:  SANTIAGO T. PEÑA JR.		Approved by:  EDGARDO E. TULIN
	END USER		UNIT HEAD, PROJECT LEADER		<i>shw</i> PRESIDENT, VSU